

For questions about the procedure call 609-265-1700

MoviPrep - Option 2 (SPLIT DOSING)

You may use this option if your colonoscopy is in the afternoon.

FIRST DOSE: Begin after Work, but no later than 7 PM. **After mixing pour the MoviPrep over a glass of ice to make it cold.** This makes the MoviPrep taste better. (Look at the Illustration below on how to mix MoviPrep)

SECOND DOSE: Begin the second dose 5 ½ hours before leaving your House to come in for your Colonoscopy. Remember pour the MoviPrep over ice to make it taste better.

Patient Instructions

The MoviPrep carton contains 4 pouches and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

For Each of the Two Doses

STEP 1	<ul style="list-style-type: none">Empty 1 Pouch A and 1 Pouch B into the disposable containerAdd lukewarm drinking water to the top line of the container. Mix to dissolve <p>If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.</p>	
STEP 2	<ul style="list-style-type: none">The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is completeDrink 16 oz of the clear liquid of your choice	

THE MORNING OF THE PROCEDURE:

AFTER FINISHING THE MORNING DOSE OF THE MOVIPREP: Do Not Eat or Drink ANYTHING More the day of the test, except morning medicines with a small sip of water up to 4 hours before the test. DO NOT CHEW GUM, HARD CANDY, or SMOKE UNTIL AFTER YOUR PROCEDURE.