

For questions about the procedure call 609-265-1700

### MoviPrep - Option 3

#### (SPLIT DOSING - Both Doses the Day before the Colonoscopy)

You may use this option if you are not working the day before the test.



**FIRST DOSE:** Begin in the morning of the day before the colonoscopy. (We suggest you mix the solution the night before) **After mixing pour the MoviPrep over a glass of ice to make it cold, if it is not cold already.** This makes the MoviPrep taste better. Once you finish the morning dose mix the solution again and place in the refrigerator to get it cold for when you drink the second dose. (Look at the Illustration below on how to mix MoviPrep)

**SECOND DOSE:** Begin in the late afternoon, but no later than 7 pm on the day before your colonoscopy. Remember pour the MoviPrep over ice to make it taste better, if it is not cold already.

## Patient Instructions

The MoviPrep carton contains 4 pouches and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

### For Each of the Two Doses

<p><b>STEP 1</b></p>	<ul style="list-style-type: none"> <li>■ Empty 1 Pouch A and 1 Pouch B into the disposable container</li> <li>■ Add lukewarm drinking water to the top line of the container. Mix to dissolve</li> </ul> <p>If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.</p>	
<p><b>STEP 2</b></p>	<ul style="list-style-type: none"> <li>■ The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is complete</li> <li>■ Drink 16 oz of the clear liquid of your choice</li> </ul>	

#### THE MORNING OF THE PROCEDURE:

**Do Not Eat or Drink ANYTHING More the day of the test, except morning medicines with a small sip of water up to 4 hours before the test. DO NOT CHEW GUM, HARD CANDY, or SMOKE UNTIL AFTER YOUR PROCEDURE.**